

CREATE[®]

300 FS

SEE ENCLOSED LEAFLET FOR FULL DETAILS

SIEN INGESLOTE PAMFLET VIR VOLLEDIGE
BESONDERHEDE

Reg. No. L10658 (Act No. 36 of 1947)

Reg. Nr. L10658 (Wet Nr. 36 van 1947)

A flowable concentrate systemic seed dressing for the control of seed decay and damping-off as well as the suppression of cob and tassel smut caused by *Sphacelotheca reiliana* in maize.

'n Vloeibare konsentraat sistemiese saadbehandelingsmiddel vir die beheer van saadverrotting en saailing-verwelksiekte asook die onderdrukking van kop- en pluimvrot wat veroorsaak word deur *Sphacelotheca reiliana* in mielies.

FUNGICIDE GROUP CODE 3 SWAMDODER GROEPKODE

Active ingredient:

Prothioconazole (triazole)..... 300 g/l

Aktiewe bestanddeel:

Prothioconazole (triazole)..... 300 g/l

REGISTERED BY / GEREGISTREER DEUR:

Bayer (Pty) Ltd / (Edms) Bpk

Reg. 1968/011192/07

P.O. Box/Posbus 143

Isando

1600 South Africa/Suid-Afrika

TEL: (OFFICE HOURS) / (KANTOORURE) (011) 921 5911

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL: 086 155 5777

Create[®] is a registered trademark of the Bayer Group.

Create[®] is 'n geregistreerde handelsmerk van die Bayer Groep.

© Bayer (Pty) Ltd

BATCH NUMBER:

See on pack

LOTNOMMER:

DATE OF MANUFACTURE:

Sien op houer

DATUM VERVAARDIG:

Bayer

RSA/0719/Create 300 FS 200L/Sales Panel
Code: 85332031A



CREATE[®]

300 FS

Seed Treatment

A flowable concentrate systemic seed dressing for the control of seed decay and damping-off as well as the suppression of cob and tassel smut caused by *Sphacelotheca reiliana* in maize.

START

Packaging / Verpakking:
200 L & 1000 L



CAUTION
VERSIGTIG



Saadbehandelingsmiddel

WAARSKUWINGS:

BEHANDEL SLEGS SAAD WAT GEPLANT GAAN WORD. MOET NOOIT BEHANDELDE SAAD AS VOEDSEL OF VOER GEBRUIK NIE.

MOENIE BINNE 85 DAE NA PLANT OES OF BEWEI.

Gebruik **Create 300 FS** slegs vir die behandeling van hoë gehalte skoongemaakte saad.

Uiters giftig vir waterlewe met langdurige effek.

Bewaar weg van voedsel en voer.

Hou buite bereik van kinders, oningeligte persone en diere.

In die geval van vergiftiging ONTBIED 'N GENEESHEER EN TOON ETIKET AAN HOM/HAAR.

Optimale saadbedtoestande is 'n vereiste vir vinnige ontkieming en opkoms van mielies. Deeglike saadbedvoorbereiding is dus noodsaaklik vir sukses. Dit is bekend dat opkoms nadelig beïnvloed mag word deur faktore soos: 'n baie droë-, buitengewoon los- en kluitrige saadbed, gronde wat neig om toe te slaan (oppervlakte kompaksie), versuip-toestande, te diep of te vlak plant-of saaidiepte, die gebruik van saad met 'n lae persentasie ontkieming en swak kiemkragtigheid ens. Indien oppervlaktekompaksie voorkom na plant en voor opkoms moet die kors gebreek word met 'n geskikte implement.

ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GETOETS IS, WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE AANGESIEN DIE WERKING EN EFFEK DAARVAN BEÏNVOED KAN WORD DEUR FAKTORE SOOS ABNORMALE GROND-, KLIMAATS- EN BERGINGSTOESTANDE; VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE EN DIE VOORKOMS VAN WEERSTAND VAN DIE SIEKTE TEEN DIE BETROKKE MIDDEL SOWEL AS DIE METODE, TYD EN AKKURAAKHEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADELIGE EFFEK OP MENS OF DIER OF VIR 'N GEBREK AAN PRESTASIE AS GEVOLG VAN DIE VERSUIM VAN DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN TOESTANDE VAN TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID.

VOORSORGMATREËLS:

Dra beskermende handskoene, beskermende klere, oogbeskerming en gesigbeskerming tydens gebruik.

Was dadelik met seep en water na hantering.

Moet nie eet, drink of rook tydens toediening of voordat hande en gesig gewas is nie.

Spoel leë houer drie keer uit met 'n volume water gelyk aan ten minste een tiende van dié van die houer en voeg die uitspoelwater by die inhoud van die mengbak voordat die houer vernietig word.

Vernietig leë houer deur gate in te maak en plat te slaan. Moet vir geen ander doel weer gebruik nie.

Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.

Hanteer behandelde saad versigtig. Merk sakke wat behandelde saad bevat duidelik "GIFTIG" en moet nooit die saadsakke vir enige ander doel gebruik nie.

WEERSTANDBESTUUR:

Vir weerstandsbestuur is **Create 300 FS** 'n groepkode 3 swamdoder. Enige swampopulasie mag individue insluit wat 'n natuurlike weerstand teen **Create 300 FS** en ander groepkode 3 swamdoders het. Indien hierdie swamdoders herhaaldelik aangewend word kan die weerstandbiedende individue uiteindelik die swampopulasie oorheers. Hierdie weerstandbiedende swamme sal waarskynlik nie deur **Create** of enige ander groepkode 3 swamdoder beheer word nie.

Vir spesifieke inligting oor weerstandsbestuur kontak die registrasiehouer van hierdie produk.



GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANBEVEEL

Saad mag slegs behandel word met behulp van industriële saadbehandelingsmasjiene. Die masjiene moet met gereelde intervalle nagegaan word om te verseker dat die korrekte hoeveelheid **Create 300 FS** aangewend word en dat saad deeglik bedek word met die swamdoder.

Meng die benodigde volume **Create 300 FS** deeglik met die benodigde volume water om 'n totale mengsel volume van 1000 ml/100 kg saad op te maak voordat die saad daarmee behandel word.

Alle opgemaakte "slurries" wat **Create 300 FS** bevat moet binne 24 uur na opmaak gebruik word.

Mieliesaad kan onmiddellik voor plant met **Create 300 FS** behandel word of vroeër en dan opgeberg word tot planttyd sonder dat ontkieming en swamdodende aktiwiteit benadeel sal word.

VERENIGBAARHEID:

Create 300 FS is verenigbaar met **Poncho** en **Poncho Votivo**.

TOEDIENINGSTABEL:

GEWAS/SIEKTE	TOEDIENINGSHOEVEELHEID	OPMERKINGS
MIELIES Saadverrotting en saailing- vewelksiekte (<i>Fusarium</i> spp.)	0.1 ml / 1000 saad	Verseker eweredige en deeglike bedekking van saad
Onderdrukking van kop-en pluimvrot (<i>Sphacelotheca</i> <i>reiliana</i>)	0.3 ml / 1000 saad	Sien GEBRUIKSAANWYSINGS

Create®, **Poncho**® en **Poncho Votivo**® is die geregistreerde handelsmerke van die Bayer Groep.

© Bayer (Pty) Ltd. All unauthorized reproduction of this label is forbidden.

Bayer

Seed Treatment

WARNINGS:

TREAT ONLY SEED THAT IS TO BE USED FOR PLANTING PURPOSES. NEVER USE TREATED SEED FOR HUMAN CONSUMPTION OR ANIMAL FEED.

DO NOT HARVEST OR GRAZE WITHIN **85** DAYS AFTER PLANTING.

Use **Create 300 FS** only for the treatment of high quality cleaned seed.

Very toxic to aquatic life with long lasting effects.

Store away from food and feedstuffs.

Keep out of reach of children, uninformed persons and animals.

In case of poisoning CALL A DOCTOR AND MAKE LABEL AVAILABLE TO HIM/HER.

Optimal seedbed and planting conditions are necessary for proper germination and emergence of maize. Thorough seedbed preparation is essential for success. It is known that reduced crop emergence can be caused by adverse factors such as: a very dry, exceptionally loose and cloddy seedbed, soils that tend to crust (surface compaction), water logging, drilling/sowing too deep or too shallow, use of seed with low germination capacity and vigor etc. Should surface compaction of the soil take place after planting and before emergence, this must be broken by means of a suitable implement.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL SOIL, CLIMATIC AND STORAGE CONDITIONS; COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE DISEASE AGAINST THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

Wear protective gloves/ protective clothing/ eye protection/ face protection.

Wash with soap and water immediately after use.

Do not eat, drink or smoke whilst applying or before washing hands and face.

Rinse empty container three times with a volume of water equal to at least one tenth of that of the container and add the rinsings to the contents of the mixing tank before disposing of the container.

Destroy empty container by perforation and flattening and never use for any other purpose.

Prevent contamination of food, feedstuffs, drinking water and eating utensils.

Handle treated seed carefully and clearly mark the bags which contain treated seed "POISONOUS" and never use the seed bags again for other purposes.

RESISTANCE MANAGEMENT:

For resistance management, **Create 300 FS** is a group code 3 fungicide. Any fungus population may contain individuals naturally resistant to **Create 300 FS** and other group code 3 fungicides. The resistant individuals can eventually dominate the fungus population if these fungicides are used repeatedly. These resistant fungi may not be controlled by **Create** or any other group code 3 fungicides.

For specific information on resistance management contact the registration holder of this product.



DIRECTIONS FOR USE: USE ONLY AS DIRECTED

Seed must only be treated by means of industrial seed dressing machines. The machines should be checked at regular intervals to ensure that the correct quantity of **Create 300 FS** is used and that the seed is thoroughly covered with the fungicide.

Mix the required volume of **Create 300 FS** thoroughly with the required volume of water to make up a slurry volume equivalent to 1000 ml/100 kg seed before treating the seed.

All slurry mixtures that contain **Create 300 FS** should be used within 24 hours of mixing.

Maize seed may be treated with **Create 300 FS** immediately before planting, but can be treated earlier and stored under cool dry conditions for a season without germination being affected.

COMPATIBILITY:

Create 300 FS is compatible with **Poncho** and **Poncho Votivo**.

APPLICATION TABLE:

CROP/DISEASE	DOSAGE	REMARKS
MAIZE Seed decay and damping-off (<i>Fusarium</i> spp.)	0.1 ml / 1000 seed	Ensure even and thorough coverage of the seed
Suppression of Cob and Tassel smut (<i>Sphacelotheca reiliana</i>)	0.3 ml / 1000 seed	See DIRECTIONS FOR USE

Create®, **Poncho**® and **Poncho Votivo**® are the registered trademarks of Bayer Group.

© Bayer (Pty) Ltd. All unauthorized reproduction of this label is forbidden.

Bayer

RSA/0719/Create 300 FS 200L/Sales Panel
Code: ZAB5332031A

