



SEE ENCLOSED PAMPHLET FOR FULL PARTICULARS

Reg. No. L8525 (Act No. 36 of 1947)
Namibian Reg No. N-AR 1572

A suspension concentrate post-emergence herbicide for the control of certain broadleaf and grass weeds in maize and sugarcane.

SIEN INGESLOTE PAMFLET VIR VOLLEDIGE BESONDERHEDE

Reg. Nr. L8525 (Wet Nr. 36 van 1947)
Namibiese Reg. Nr. N-AR 1572

'n Suspensie konsentraat na opkoms onkruidodder vir die beheer van sekere breëblaar- en graskonkruid in mielies en suikerriet.

HERBICIDE GROUP CODE F₂ ONKRUIDDODER GROEPKODE

Active ingredients:

Tembotrione (Triketone).....420 g/l
Isoxadifen-ethyl (Safener).....210 g/l

Aktiewe bestanddele:

Tembotrione (Triketone).....420 g/l
Isoxadifen-etiel (Beveiliging).....210 g/l

REGISTERED BY / GEREGISTREER DEUR:

Bayer (Pty) Ltd / (Edms) Bpk
Reg. 1968/011192/07
P.O. Box/Posbus 143
Isando
1600 South Africa/Suid-Afrika
TEL: (OFFICE HOURS) / (KANTOORURE) (011) 921 5911

IN CASE OF POISONING PLEASE PHONE / IN GEVAL VAN VERGIFTIGING SKAKEL: 0861555777

Laudis® is a registered trademark of the Bayer Group.

Laudis® is 'n geregistreerde handelsmerk van die Bayer Groep.

© Bayer (Pty) Ltd

BATCH NUMBER: See on pack
LOTNOMMER:

DATE OF MANUFACTURE: Sien op houer
DATUM VERVAARDIG:

Bayer


RSA/0318/Laudis 900ml/Sales Panel
Code: 84951498D



Herbicide

A suspension concentrate post-emergence herbicide for the control of certain broadleaf and grass weeds in maize and sugarcane.

contains:
Tembotrione (Triketone) and
Isoxadifen-ethyl (Safener)



**Packaging / Verpakking:
900 ml & 5 L**



**HARMFUL
SKADELIK**



Onkruidodder

WAARSKUWINGS:

MOET NIE BEWEI OF PLANTE VAN BEHANDELDE MIELIES AS VOER GEBRUIK BINNE 70 DAE NA LAASTE TOEDIENING NIE.

MOET NIE BEWEI OF PLANTE VAN BEHANDELDE SUIKERRIET AS VOER GEBRUIK BINNE 235 DAE NA LAASTE TOEDIENING NIE.

Hanteer versigtig.

Giftig wanneer ingesluk.

Kan vermoedelik skadelik wees vir die ongebore kind.

Mag skade aan organe (oë, niere, lewer) veroorsaak deur langdurige en herhaaldelike blootstelling.

Baie giftig vir water lewe met langdurige effek.

Berg in 'n koel plek, weg van voedsel, voer, saad of enige ander landboumiddels.

Hou buite bereik van kinders, oningeligte persone en diere.

Voorkom kontak met oë, vel en klere.

Herbetreding: Moenie behandelde gebied binne een dag na toediening binnegaan nie, tensy beskermende oorklere gedra word.

Laudis is op die meeste belangrike kultivars getoets en geen betekenisvolle simptome van fitotoksiteit in die vorm van oesverlagings het onder normale groeitoestande voorgekom nie. **Laudis** moet nie toegedien word op mielies of suikerriet wat skade simptome van vorige onkruidodder toediening vertoon nie.

ALHOEWEL HIERDIE MIDDEL OMVATTEND ONDER 'N GROOT VERSKEIDENHEID TOESTANDE GETOETS IS WAARBORG DIE REGISTRASIEHOUER NIE DAT DIT ONDER ALLE TOESTANDE DOELTREFFEND SAL WEES NIE AANGESIEN DIE WERKING EN EFFEK DAARVAN BEINVLOED KAN WORD DEUR FAKTORE SOOS ABNORMALE GROND-, KLIMAATS- EN BERGINGSTOESTANDE, KWALITEIT VAN VERDUNNINGSWATER, VERENIGBAARHEID MET ANDER STOWWE WAT NIE OP DIE ETIKET AANGEDUI IS NIE EN DIE VOORKOMS VAN WEERSTAND VAN DIE ONKRUIDE TEEN DIE BETROKKE MIDDEL SOWEL AS DIE METODE, TYD EN AKKURAAKHEID VAN TOEDIENING. VERDER AANVAAR DIE REGISTRASIEHOUER NIE VERANTWOORDELIKHEID VIR SKADE AAN GEWASSE, PLANTEGROEI, DIE OMGEWING OF VIR NADELIGE EFFEK OP MENS OF DIER OF VIR 'N GEBREK AAN PRESTASIE VAN DIE BETROKKE MIDDEL AS GEVOLG VAN DIE VERSUIM VAN DIE GEBRUIKER OM ETIKETAANWYSINGS NA TE KOM OF AS GEVOLG VAN DIE ONTSTAAN VAN DIE TOESTANDE WAT NIE KRAGTENS DIE REGISTRASIE VOORSIEN KON WORD NIE. RAADPLEEG DIE VERSKAFFER IN DIE GEVAL VAN ENIGE ONSEKERHEID.

VOORSORGMAATREËLS:

Verhoed inaseming van spuitdampe.

Dra beskermende klere (oorpak, stewels, rubberhandskoene) wanneer die produk hanteer of toegedien word.

Was met seep en water onmiddellik na gebruik of toevallige velkontak.

Moenie eet, drink of rook tydens vermenging, toediening of voordat hande en gesig gewas is nie.

Was besoedelde klere daagliks.

Voorkom oorwaai van spuitstof na ander gewasse, weiding, riviere en damme.

Maak spuitapparaat skoon na gebruik, raak ontslae van uitspoelwater waar dit nie gewasse, weiding, riviere en damme kan besoedel nie.

Spoel leë houer drie keer uit met 'n volume water gelyk aan ten minste een tiende van die houer en voeg die uitspoelwater by die inhoud van die spuittenk voordat van die houer op die voorgeskrewe metode ontslae geraak word.

Vernietig leë houer deur gate daarin te maak en plat te slaan. Moet vir geen ander doel weer gebruik nie.

Voorkom besoedeling van voedsel, voer, drinkwater en eetgerei.



GEBRUIKSBEPERKINGS:

1. Om skade op opvolggewasse te voorkom, verwys na die wagperiodes op die **Atrazine 500 SC** etiket.
2. Vermoed dubbele bespuiting en die oormatige oorvlueing van spuitbane.
3. Oormatige toediening, toediening tydens die verkeerde groeistadium, of enige ander gebruik wat nie in ooreenstemming met hierdie etiket as gebruiksaanwysings is nie, kan gewasskade tot gevolg hê.
4. **Laudis** is nie getoets op ingeteelde oerplante van basters, eksperimentele basters of kultivars, nuut vrygestelde basters of kultivars, suikermielies of springmielies nie. **Moet nie Laudis** toedien alvorens 'n verteenwoordiger van Bayer CropScience of saad voorsieners nie geraadpleeg is nie.

SIMPTOME VAN VERGIFTIGING:

Skadelik indien ingesluk. Mag sensitiwiteit van die vel veroorsaak. Mag ligte oog-irritasie veroorsaak.

NOODHULPBEHANDELING:

Inaseming: Verskuif na vars lug. Kontak onmiddellik 'n geneesheer of die Gif Beheer Sentrum.

Velkontak: Was deeglik met seep en genoegsame water en indien beskikbaar met poli-etileenglykool 400 en spoel af met water.

Oogkontak: Spoel onmiddellik vir ten minste 15 minute met genoegsame water, ook onder die ooglede. Verwyder kontaklense na die eerste 5 minute indien teenwoordig en hou aan met spoeling.

Ingesluk: Braking MOET NIE veroorsaak word nie. Kontak onmiddellik 'n geneesheer of die Gif Beheer Sentrum. Spoel mond uit.

NOTA AAN GENEESHEER:

Behandeling: Wanneer gif ingesluk is, doen maagspoeling indien nodig.

WEERSTANDSWAARSKUWING:

Laudis behoort aan die Trikitone groep van onkruidodders. Die metode van werking van hierdie groep is die inhibering van die 4-HPPD ensiem en as sulks word dit deur HRAC as 'n Groepkode F₂ onkruidodder geklassifiseer. Enige populasie van 'n spesifieke onkruid mag individue insluit wat 'n natuurlike weerstand teen **Laudis**, of enige ander Groepkode F₂ onkruidodder het. Indien hierdie onkruidodders herhaaldelik aangewend word, kan die weerstandbiedende individue uiteindelik die onkruidpopulasie oorheers. Hierdie weerstandbiedende onkruid sal moontlik nie deur **Laudis** of enige ander Groepkode F₂ onkruidodder, beheer word nie.

Onkruidodder

Aangesien die voorkoms van weerstandbiedende onkruid moeilik is om te bepaal voor onkruidodertoediening plaasvind, is dit uiters noodsaaklik dat behandelde gebiede met gereelde intervalle ondersoek word om weerstand vroegtydig te identifiseer. Bayer (Edms) Bpk. sal nie verantwoordelikheid aanvaar vir enige gebrek aan prestasie in die geval van die opbou van weerstand in die onkruidpopulasie as gevolg van onvoldoende onkruidbestandheidsbestuur soos aangedui op die etiket nie.

Om weerstand teen onkruidodders te vertraag:

- Vermyn die eksklusiewe, herhaaldelike gebruik van onkruidodders in dieselfde onkruidodder groepkode.
- Wissel af met produkte in verskillende onkruidodder groepkodes.
- Integreer ander beheermaatreëls (chemies, verbouing, ens.) in onkruidbeheerprogramme.

Vir spesifieke inligting oor weerstandsbestuur, kontak u naaste Bayer CropScience verteenwoordiger.

GEBRUIKSAANWYSINGS: GEBRUIK SLEGS SOOS AANGEDUI

VERENIGBAARHEID:

Laudis is verenigbaar met insekdoders **Bulldock Beta 125 SC** (Reg. Nr. L7612), **Decis Forte** (Reg. Nr. L6563), **Larvin 375 SC** (Reg. Nr. L2997); en onkruidodders **Harness Extra EC** (Reg. Nr. L7703), **Mamba DMA 480 SL** (Reg. Nr. L8388) en **Roundup PowerMax** (Reg. Nr. L7769). **Laudis** moet nie gemeng word met enige produkte, anders as dié wat hierbo genoem word en waarna verwys word in die TOEDIENINGSTABEL nie.

MENG- EN TOEDIENINGSINSTRUKSIES:

GEBRUIK DIE SPUITMENGSEL BINNE 6 UUR NA VERMENGING

1. Maak 'n vooraf mengsel in 'n aparte houër met die aanbevole hoeveelhede van produkte in die volgende volgorde: **Laudis**, **Atrazine 500 SC** (+ insekdoder + **Harness Extra** – indien nodig) en dan laaste **Ballista** in ongeveer 10 liters skoon water.
2. Maak die spuittenk halfvol met water.
3. Voeg dan die aanbevole hoeveelheid **ammonium sulfaat** by. Verseker dat dit volledig opgelos is voordat die voorafgemengde mengsel bygevoeg word.
4. Voeg die voorafgemengde mengsel (verwys na punt 1 hierbo) in die spuittenk en vul die tenk verder met water.
5. Vir meng instruksies (volgorde, ens.) met **Roundup PowerMax** of **Mamba DMA 480 SL**, verwys asseblief na die onderskeie etikette.
6. Hou die mengsel in die spuittenk voortdurend in suspensie deur middel van roering.
7. Verseker eweredige bedekking van die onkruidblare deur gebruik te maak van spuitpunte wat geskik is vir optimale na-opkoms toedienings, soos byvoorbeeld dubbel uitlaat waaiertipe (Twinjet) spuitpunte.
8. Dien toe by wyse van 'n trekker-gemonteerde balkspuit, met volgehoue roering en wat korrek gekalibreer is om ten minste 200 - 300 l spuitmengsel/ha toe te dien.
9. Dien so gou moontlik na vermenging toe vir optimale resultate.
10. Aan die einde van elke dag se bespuiting moet die spuittenk deeglik met water uitgespoel word.



TOEDIENINGSTABEL:

GEWAS	DOSIS PER HEKTAAR	OPMERKINGS
MIELIES (Nie vir gebruik op suikermielies)	Laudis - 120 ml + Ballista - 500 ml + Atrazine 500 SC - 1000 ml + Ammonium sulfaat (AMS) 2% Oplossing indien 'n 500 g/l vloeibare formulاسie van AMS gebruik word. OF 1% oplossings indien 'n 1000 g/kg korrel formulاسie van AMS gebruik word.	Dien toe wanneer onkruid in die 4 tot 6-blaar groeistadium is. Verseker totale bedekking van onkruidblare. Bereken die persentasie AMS volgens die watervolume wat per hektaar toegedien word.

OF

GEWAS	DOSIS PER HEKTAAR	OPMERKINGS
MIELIES (Nie vir gebruik op suikermielies)	Laudis - 180 ml + Ballista - 500 ml + Atrazine 500 SC - 1000 ml	Dien toe wanneer onkruid in die 4 tot 6-blaar groeistadium is. Verseker totale bedekking van onkruidblare. Verwys asseblief na die beperking op onkruidspesies aangedui in onderstaande lys wat gemerk is met * [1] [2].

Onkruidodder



GLIFOSAAT-VERDRAAGSAME MIELIES

GEWAS	DOSIS PER HEKTAAR	OPMERKINGS
Glifosaat verdraagsame mielies (Uitsluitend suikermielies)	Laudis - 180 mℓ + Roundup Powermax OF Mamba DMA 480 SL + Ammonium sulphate (AMS)	Vir verdere besonderhede oor onkruidbeheer en toedienings instruksies (water volume, meng instruksies, ens.) raadpleeg asseblief die onderskeie Roundup PowerMax en Mamba DMA 480 SL etikette.
SUIKERRIET	Laudis - 180 mℓ + Ballista - 500 mℓ + Atrazine 500 SC - 1000 mℓ + Ammonium sulfaat (AMS) 2% Oplossing indien 'n 500 g/ℓ vloeibare formulاسie van AMS gebruik word. OF 1% oplossings indien 'n 1000 g/kg korrel formulاسie van AMS gebruik word.	Dien toe wanneer onkruid in die 4 tot 6-blaar groeistadium is en aktief groei. Die beste resultate sal verkry word gedurende warm en vogtig groei toestande. Verseker totale bedekking van onkruidblare. Bereken die persentasie AMS volgens die watervolume wat per hektaar toegedien word.

BREëBLAARONKRUIDE BEHEER:

<i>Acanthospermum hispidum</i>	Regopsterklits
<i>Ageratum conyzoides</i>	Bokkruid
<i>Amaranthus hybridus</i>	Gewone misbredie
<i>Amaranthus deflexus</i>	Meerjarige misbredie
<i>Amaranthus retroflexus</i>	
<i>Amaranthus spinosus</i>	Doringmisbredie
* [1] <i>Bidens pilosa</i>	Gewone knapsekêrel
<i>Chenopodium album</i>	Withondebossie
<i>Chenopodium carinatum</i>	Groenhondebossie
<i>Chenopodium pumilio</i>	
<i>Cleome monophylla</i>	Rusperbossie (enkelblaar cleome)
<i>Commelina benghalensis</i>	Bengaalse wandelende Jood
<i>Corchorus olitorius</i>	Indiese jute
<i>Datura ferox</i>	Grootstinkblaar
<i>Datura stramonium</i>	Olieboom
<i>Galinsoga parviflora</i>	Knopkruid
<i>pomoea purpurea</i>	Purperwinde
<i>Ipomoea obscura</i>	Wildepatat
<i>Nicandra physaloides</i>	Basterappelliefie
<i>Physalis angulata</i>	Wilde-appelliefie
<i>Portulaca oleracea</i>	Porslein
<i>Richardia brachiensis</i>	Meksikaanse richardia
<i>Schkura pinnata</i>	Kleinkakiebos
<i>Tagetes minuta</i>	Kakiebos
<i>Tribulus terrestris</i>	Dubbeltjie
<i>Galinsoga parviflora</i>	Knopkruid
<i>Xanthium strumarium</i>	Kankerroos

GRASONKRUIDE BEHEER:

<i>Crotalaria sphaerocarpa</i>	Wildeluserm
* [2] <i>Dactyloctenium aegyptium</i>	Hoenderspoor
<i>Digitaria sanguinalis</i>	Kruisingergras
<i>Echinochloa colona</i>	Moerasgras
<i>Eleusine coracana</i>	Jongosgras
<i>Panicum maximum</i>	Gewone buffelsgras
<i>Paspalum dilatatum</i>	Gewone Paspalum
<i>Sorghum verticilliflorum</i>	Gewone wildesorghum
<i>Urochloa mosambicensis</i>	Bosveldbeesgras
<i>Urochloa panicoides</i>	Beesgras

*[1] [2] Slegs **onkruidonderdrukking** indien **Laudis** toegedien word teen 180 mℓ/ha **sonder** Ammonium sulfaat. Vir **goeie beheer** verseker dat Ammonium Sulfaat **bygevoeg** word.

NOTA:

- Laudis** beheer bogenoemde onkruidspesies. Ander onkruidspesies wat nie tydens die ontwikkelingsproewe met die middel voorgekom het nie, kan moontlik in 'n meerdere of mindere mate beheer word. Die registrasiehouer aanvaar egter geen aanspreeklikheid ten opsigte van effektiwiteit op ongelyste onkruid nie.
- Moet nie **Laudis** toedien wanneer Mielies of onkruid onder stremming (koue, droogte, ens.) is nie.
- Laudis** mag chlorose van mielieblare veroorsaak. Hierdie simptome is egter tydelik van aard en opbrengs sal nie beïnvloed word nie.
- Laudis** mag chlorose en verdwering van suikerriet plante veroorsaak. Hierdie simptome is van 'n verbygaande aard en sal nie opbrengs beïnvloed nie.
- Die aanbevole gebruik van **Laudis** in 'n onkruidodder spuitprogram is as volg:
 - Tydens plant – Dit word aanbeveel dat n vooropkoms onkruidbeheer program gevolg word.
 - Ongeveer 4-6 weke later (afhange van die onkruid groeistadium) dien 'n algehele bespuiting van **Laudis** toe soos aanbeveel in die TOEDIENINGSTABEL.
- Vir nawerkende gras beheer in mielies kan **Harness Extra EC** by die geregistreerde **Laudis** mengsel gevoeg word. Raadpleeg die **Harness Extra** etiket vir meer inligting oor die maontlike spektrum wat beheer kan word met so mengsel.

Larvin®, **Bulldock®** en **Decis®** is geregistreerde handelsmerke van die Bayer Groep.
Roundup PowerMAX® en **Harness®** is geregistreerde handelsmerke van Monsanto Suid Afrika (Edms), Bpk.
Mamba® is geregistreerde handelsmerk van Dow AgroSciences Suidelike Afrika (Edms) Bpk.

© Bayer (Edms) Bpk. Alle ongemagtigde reproduksie van hierdie etiket word verbied.

Herbicide

WARNINGS:

DO NOT GRAZE OR USE PLANTS AS FODDER FROM TREATED MAIZE WITHIN 70 DAYS OF LAST APPLICATION.

DO NOT USE PLANTS AS FODDER FROM TREATED SUGARCANE WITHIN 235 DAYS OF LAST APPLICATION.

Handle with care.

Harmful if swallowed.

Suspected of damaging the unborn child.

May cause damage to organs (eyes, kidneys, liver) through prolonged or repeated exposure.

Very toxic to aquatic life with long lasting effects.

Store in a cool place away from food, feed, seed and other agricultural remedies.

Keep out of reach of children, uninformed persons and animals.

Avoid contact with eyes, skin or clothing.

RE-ENTRY: Do not enter treated area within one day after treatment unless wearing protective clothing.

Laudis has been tested on most important cultivars and no significant phytotoxicity in the form of yield reduction has been recorded under normal growing conditions. Do not apply **Laudis** to maize or sugarcane that exhibits injury from previous herbicide applications.

ALTHOUGH THIS REMEDY HAS BEEN EXTENSIVELY TESTED UNDER A LARGE VARIETY OF CONDITIONS THE REGISTRATION HOLDER DOES NOT WARRANT THAT IT WILL BE EFFICACIOUS UNDER ALL CONDITIONS BECAUSE THE ACTION AND EFFECT THEREOF MAY BE AFFECTED BY FACTORS SUCH AS ABNORMAL SOIL, CLIMATIC AND STORAGE CONDITIONS, QUALITY OF DILUTION WATER, COMPATIBILITY WITH OTHER SUBSTANCES NOT INDICATED ON THE LABEL AND THE OCCURRENCE OF RESISTANCE OF THE WEEDS TO THE REMEDY CONCERNED AS WELL AS BY THE METHOD, TIME AND ACCURACY OF APPLICATION. THE REGISTRATION HOLDER FURTHERMORE DOES NOT ACCEPT RESPONSIBILITY FOR DAMAGE TO CROPS, VEGETATION, THE ENVIRONMENT OR HARM TO MAN OR ANIMAL OR FOR LACK OF PERFORMANCE OF THE REMEDY CONCERNED DUE TO FAILURE OF THE USER TO FOLLOW THE LABEL INSTRUCTIONS OR TO THE OCCURRENCE OF CONDITIONS WHICH COULD NOT HAVE BEEN FORESEEN IN TERMS OF THE REGISTRATION. CONSULT THE SUPPLIER IN THE EVENT OF ANY UNCERTAINTY.

PRECAUTIONS:

Do not inhale spray mist.

Wear protective clothing (overall, boots, rubber gloves) when handling or applying the product.

Wash with soap and water immediately after use or accidental skin contact.

Do not eat, drink or smoke whilst mixing, applying or before washing hands and face.

Wash contaminated clothing daily.

Avoid spray drift onto other crops, grazing, rivers and dams.

Clean applicator after use, dispose of wash water where it will not contaminate crops, grazing, rivers and dams.

Rinse empty container three times with a volume water equal to at least one tenth of that of the container and add the rinsing to the contents of the spray tank before disposing of the container in the prescribed manner.

Destroy empty container by perforation and flattening and never use for any other purpose.

Prevent contamination of food, feedstuffs, drinking water and eating utensils.

USE RESTRICTIONS:

1. To avoid damage to follow-up crops, refer to the waiting periods on the **Atrazine 500 SC** label.
2. Avoid excessive overlapping of spray swaths and double spraying.



3. Over-application, application at the incorrect growth stage, or any other use not in accordance with the directions on this label may cause crop damage.
4. **Laudis** has not been tested on inbred parent plants of maize hybrids, experimental hybrids or cultivars, newly released hybrids or cultivars, sweetcorn or popcorn. **Do not** apply **Laudis** without first consulting a representative of Bayer CropScience or the seed supplier.

SYMPTOMS OF POISONING:

Harmful if swallowed. May cause sensitization by skin contact. May cause mild eye irritation.

FIRST AID TREATMENT:

Inhalation: Move to fresh air. Call a physician or poison control center immediately.

Skin contact: Wash off thoroughly with plenty of water and soap, if available with polyethyleneglycol 400, subsequently rinse with water.

Eye contact: Rinse immediately with plenty of water, also under the eyelids, for at least 15 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

Ingestion: Do NOT induce vomiting. Call a physician or poison control center immediately. Rinse mouth.

NOTE TO PHYSICIAN:

Treatment: When swallowed, if necessary flush stomach.

RESISTANCE WARNING:

Laudis belongs to the Triketone group of herbicides. The mode of action of this group is the inhibition of the 4-HPPD enzyme and, as such it is classified by HRAC as a Group Code F₂ herbicide. Any weed population may contain individuals naturally resistant to **Laudis** and other Group Code F₂ herbicides. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds may not be controlled by **Laudis** or any other Group Code F₂ herbicides.

Herbicide

Since the presence of resistant weeds is difficult to detect prior to herbicide application, it is of the utmost importance that treated areas be inspected at regular intervals to identify the occurrence of herbicide-resistant weeds timeously. Bayer (Pty) Ltd will not accept liability for performance failures in the event of a build-up of resistant weeds resulting from inadequate resistance management practices as indicated on the label.

To delay herbicide resistance:

- Avoid exclusive, repeated use of herbicides from the same herbicide Group Code.
- Alternate with products from different herbicide Group Codes.
- Integrate other control methods [chemical, cultural, etc.] in weed control programmes.

For specific information on resistance management contact your local Bayer CropScience representative.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED

COMPATIBILITY:

Laudis is compatible with insecticides **Bulldock Beta 125 SC** (Reg. No. L7612), **Decis Forte** (Reg. No. L6563), **Larvin 375 SC** (Reg. No. L2997); and herbicides **Harness Extra EC** (Reg. No. L7703), **Mamba DMA 480 SL** (Reg. No. L8388) and **Roundup PowerMax** (Reg. No. L7769).

Do not mix **Laudis** with any products, other than those mentioned above and shown in the APPLICATION TABLE.

MIXING AND APPLICATION INSTRUCTIONS:

USE THE SPRAY MIXTURE WITHIN 6 HOURS AFTER MIXING

1. Prepare a pre-mix in a separate container with the required amount of products in the following order: **Laudis**, **Atrazine 500 SC** (+ insecticide + **Harness Extra EC** – if required) and then finally **Ballista** in approximately 10 litres of clean water.
2. Half fill the spray tank with water.
3. Add the required amount of **ammonium sulphate** to the tank. Ensure that it is fully dissolved before adding the pre-mix.
4. Add the pre-mix (referred to point 1 above) to the tank and top up with water.
5. For mixing instructions (following order, etc.) with **Roundup PowerMax** or **Mamba DMA 480 SL**, please refer to the respective labels.
6. Keep the mixture in continuous suspension by means of agitation.
7. Ensure even coverage of the weed foliage by using nozzle types that are suitable for optimum post emergence application, such as double outlet flat fan (Twinjet) nozzles.
8. Apply by means of a tractor mounted boom sprayer with continual agitation and which is correctly calibrated to deliver at least 200 - 300 ℓ spray mixture/ha.
9. Spray as soon as possible after mixing for optimum results.
10. After each day's spraying, flush the tank out thoroughly with water.



APPLICATION TABLE:

CROP	APPLICATION RATE PER HECTARE	REMARKS
MAIZE (Excluding sweet corn)	Laudis - 120 ml + Ballista - 500 ml + Atrazine 500 SC - 1000 ml + Ammonium sulphate (AMS) 2% Solution when a 500 g/l flowable formulation of AMS is used. OR 1% solution when a 1000 g/kg granular formulation of AMS is used.	Apply when the weeds are between the 4 and 6 leaf growth stage. Ensure total coverage of the weed foliage. Calculate percentage AMS according to water volume applied per hectare.

OR

CROP	APPLICATION RATE PER HECTARE	REMARKS
MAIZE (Excluding sweet corn)	Laudis - 180 ml + Ballista - 500 ml + Atrazine 500 SC - 1000 ml	Apply when the weeds are between the 4 and 6 leaf growth stage. Ensure total coverage of the weed foliage. Please refer to the limitation to weed species indicated listed below marked with *(1) (2).

Herbicide

GLYPHOSATE-TOLERANT MAIZE

CROP	APPLICATION RATE PER HECTARE	REMARKS
Glyphosate tolerant maize (Excluding sweet corn)	Laudis - 180 ml + Roundup Powermax OR Mamba DMA 480 SL + Ammonium sulphate (AMS)	For further detailed on weeds controlled and application instructions (water volume, mixing instructions, etc.) please refer to the respective Roundup PowerMax and Mamba DMA 480 SL labels.
SUGARCANE	Laudis - 180 ml + Ballista - 500 ml + Atrazine 500 SC - 1000 ml + Ammonium sulphate (AMS) 2% Solution when a 500 g/l flowable formulation of AMS is used. OR 1% solution when a 1000 g/kg granular formulation of AMS is used.	Apply when the weeds are between the 4 and 6 leaf growth stage and are actively growing. Best weed control will be achieved in warm, moist growing conditions. Ensure total coverage of the weed foliage. Calculate percentage AMS according to water volume applied per hectare.

BROADLEAF WEEDS CONTROLLED:

<i>Acanthospermum hispidum</i>	Upright starbur
<i>Ageratum conyzoides</i>	Billy-goat weed
<i>Amaranthus hybridus</i>	Common pigweed
<i>Amaranthus deflexus</i>	Perennial pigweed
<i>Amaranthus retroflexus</i>	Redroot pigweed
<i>Amaranthus spinosus</i>	Spiny pigweed
* [1] <i>Bidens pilosa</i>	Common blackjack
<i>Chenopodium album</i>	White goosefoot
<i>Chenopodium carinatum</i>	Green goosefoot
<i>Chenopodium pumilio</i>	
<i>Cleome monophylla</i>	Spindlepod
<i>Commelina benghalensis</i>	Bengal wandering Jew
<i>Corchorus olitorius</i>	Indian jute
<i>Datura ferax</i>	Large thorn apple
<i>Datura stramonium</i>	Thorn apple
<i>Galisonga parviflora</i>	Gallant soldier
<i>Ipomoea obscura</i>	Small white morning glory
<i>Ipomoea purpurea</i>	Morning glory
<i>Nicandra physaloides</i>	Apple of Peru
<i>Physalis angulata</i>	Wild gooseberry
<i>Portulaca oleracea</i>	Purslane
<i>Richardia brachiensis</i>	Mexican richardia
<i>Schkuria pinnata</i>	Dwarf marigold
<i>Tagetes minuta</i>	Khaki weed
<i>Tribulus terrestris</i>	Devil's thorn
<i>Galinsoga parviflora</i>	Small-flowered quickweed
<i>Xanthium strumarium</i>	Large Cocklebur

GRASS WEEDS CONTROLLED:

<i>Crotalaria sphaerocarpa</i>	Mealie crotalaria
* [2] <i>Dactyloctenium aegyptium</i>	Crowfoot
<i>Digitaria sanguinalis</i>	Crab finger-grass
<i>Echinochloa colona</i>	Marsh grass
<i>Eleusine coracana</i>	Goose grass
<i>Panicum maximum</i>	Buffalo grass
<i>Paspalum dilatatum</i>	Common Paspalum
<i>Sorghum verticilliflorum</i>	Common wild sorghum
<i>Urochloa mosambicensis</i>	Bushveld herringbone grass
<i>Urochloa panicoides</i>	Herringbone grass

*[1] (2) Weed **suppression** only when **Laudis** is applied at 180 ml/ha **without** Ammonium sulphate. For **good control** ensure that Ammonium Sulphate is **added**.

NOTE:

- Laudis** controls the abovementioned weed species. Other weed species that were not present during the development trials with the product, may also be controlled to a certain degree. The registration holder does not accept any responsibility for efficacy on unlisted weeds.
- Do not apply **Laudis** when maize or weeds are under stress (cold, drought, etc.)
- Laudis** may cause chlorosis of maize leaves. These symptoms are of a transitory nature and yield will not be affected.
- Laudis** may cause a chlorosis and stunting of sugarcane plants. These symptoms are of a transitory nature and yield will not be affected.
- The recommended use of **Laudis** in a weed control spray program in maize is as follows:
 - At plant – It is recommended to follow a pre-emergence herbicide program.
 - Approximately 4-6 weeks later (depending on the weed growth stage) apply broadcast spray of **Laudis** as recommended in the APPLICATION TABLE.
- Harness Extra EC** can be added to the **Laudis** mixtures recommended for use in maize to achieve some residual grass control. For the grass species to be controlled by such a mixture please consult the **Harness Extra EC** label.

Larvin®, **Bulldock**® and **Decis**® are registered trademarks of the Bayer Group.
Roundup PowerMAX® and **Harness**® are registered trademarks of Monsanto South Africa (Pty), Ltd.
Mamba® is the registered trademark of Dow AgroSciences Southern Africa (Pty) Ltd.

© Bayer (Pty) Ltd. All unauthorised reproduction of this label is forbidden.



